



## Product Datasheet

<b>Product Name</b>	Visfatin Recombinant Human, His Tag
<b>Cata No</b>	CB501297
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	PBEF, Pre-B cell colony-enhancing factor, Nicotinamide phosphoribosyltransferase NAmPRTase, Nampt, MGC117256, DKFZP666B131, 1110035O14Rik.

### Description

Excess adiposity is the most important risk in the development of insulin resistance and type 2 diabetes mellitus (T2DM). Adipose tissue produces several proteins (adipocytokines) such as leptin, adiponectin, resistin, tumor necrosis factor- $\alpha$ , and IL-6, that modulate insulin sensitivity and appear to play an important role in the pathogenesis of insulin resistance, diabetes, dyslipidemia, inflammation, and atherosclerosis. However, the mechanisms by which fat tissue induces insulin resistance and the role of adipocytokines in the pathogenesis of T2DM have not been well established.

Visfatin, also known as pre-B cell colony-enhancing factor (PBEF), is a cytokine that is highly expressed in visceral fat and was originally isolated as a secreted factor that synergizes with IL-7 and stem cell factors to promote the growth of B cell precursors. Visfatin homologs have been identified in carp, invertebrate mollusks, and bacteria, as well as in vertebrates, including humans and the mouse. It has been postulated to play a role in innate immunity.

Visfatin exerts insulin-mimetic effects that are dose-dependent and quantitatively similar to those of insulin in stimulating muscle and adipocyte glucose transport, and in inhibiting hepatocyte glucose production. Intravenous injection of recombinant visfatin in mice decreased plasma glucose in a dose-dependent fashion. In keeping with its insulin-mimetic effects, visfatin was as

effective as insulin in reducing hyperglycemia in insulin-deficient diabetic mice. Visfatin was also found to be bound to and activate insulin receptor, causing receptor phosphorylation and the activation of downstream signaling molecules. However, visfatin and insulin did not compete for binding to the insulin receptor, indicating that the two proteins were recognized by different regions of the receptor. Thus, visfatin might play a role in glucose homeostasis and dysregulation in biosynthesis or signal transduction, and might contribute to the pathogenesis of diabetes.

Visfatin Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 511 amino acids and having a molecular mass of 57 kDa.

The recombinant human Visfatin is fused to His tag at N-Terminus.

### Physical Appearance

Sterile Filtered solution at a concentration of 1mg/ml.

### Purity

Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

### Formulation

Visfatin His tag protein contains 20mM Tris pH-8,

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204

Coachella, CA 92236 USA

Phone : +1.6268339877

Email : [info@cali-bio.com](mailto:info@cali-bio.com)

0.1mM DTT & 10% glycerol

## Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. **Please avoid freeze thaw cycles.**

## Sequence

MGSSHHHHHH SSGLVPRGSH MNPAAEAEFN  
ILLATDSYKV THYKQYPPNT SKVYSYFECR  
EKKTENSKLR KVKYEETVfy GLQYILNKYL  
KGVVVTKEKI QEAKDVYKEH FQDDVFNEKG  
WNYILEKYDG HLPiEiKAVP EGFVIPRGNV

## Product Datasheet

LFTVENTDPE CYWLTNWIET ILVQSWYPIT  
VATNSREQKK ILAKYLLETS GNLDGLEYKL  
HDFGYRGVSS QETAGIGASA HLVNFKGTDT  
VAGLALIKKY YGTDKDPVPGY SVPAAEHSTI  
TAWGKDHEKD AFEHIVTQFS SVPVSVVSDS  
YDIYNACEKI WGEDLRHLIV SRSTQAPLII  
RPDSGNPLDT VLKVL EILGK KFPVTENSKG  
YKLLPPYLRV IQGDGVDINT LQEIVEGMKQ  
KMWSIENIAF GSGGGLLQKL TRDLLNCSFK  
CSYVVTNGLG INVFKDPVAD PNKRSKKGRL  
SLHRTPAGNF VTLEEGKGD L EEYGQDLLHT  
VFKNGKVTKS YSFDEIRKNA QLNIELEAAH H

**\* For Non-Clinical Research Use Only \***